

Received Date : 16-May-2016
Revised Date : 11-Jul-2016
Accepted Date : 27-Jul-2016
Article type : Special Article

A collaborative study of the current concepts and challenges for treatment of anastomotic leakage in colorectal surgery

A Vallance¹, S Wexner², M. Berho², R Cahill³, M. Coleman⁴, N. Haboubi⁵, R.J. Heald⁶, R.H. Kennedy⁷, B. Moran⁶, N. Mortensen⁸, R.W. Motson⁹, R. Novell¹⁰, P.R. O'Connell¹¹, F. Ris¹², T. Rockall¹³, A. Senapati¹⁴, A. Windsor¹⁵, D.G. Jayne¹⁶

1. Royal College of Surgeons of England, London, UK
2. Cleveland Clinic, Weston, USA
3. University College Dublin, Dublin, Ireland
4. Derriford Hospital, Plymouth, UK
5. University Hospital of South Manchester, Manchester, UK
6. Basingstoke and North Hampshire Hospital, Basingstoke, UK
7. St Mark's Hospital, London, UK
8. Oxford University Hospitals, Oxford, UK
9. The ICENI Centre, Colchester University Hospital, Colchester, UK
10. The Royal Free Hospital, London, UK
11. University College Dublin, Dublin, Ireland
12. Geneva University Hospitals and Medical School, Geneva, Switzerland.
13. Royal Surrey County Hospital, Guildford, UK
14. Portsmouth Hospitals, Portsmouth, UK
15. University College Hospital, London, UK
16. University of Leeds, Leeds, UK

Corresponding author: S Wexner wexners@ccf.org (Twitter: @SWexner)

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/codi.13534

This article is protected by copyright. All rights reserved.

Disclosures: this manuscript is the outcome from a meeting of expert colorectal surgeons and pathologists held in 2016 and supported by educational grants from Novadaq Technologies Inc. (Ontario, Canada) and Elemental Healthcare (Berkshire, UK). The meeting sponsors did not participate in any aspect of planning or delivery of the academic content of the meeting nor in the drafting or critical review of the subsequent manuscript.

Abstract

The reduction of the incidence, detection, and treatment of anastomotic leakage (AL) continues to challenge the colorectal surgical community. AL is not consistently defined and reported in clinical studies, its occurrence is variably reported and its impact on long-term morbidity and healthcare resources has received relatively little attention. Controversy continues about the best strategies to reduce the risk. Diagnostic tests lack sensitivity and specificity, resulting in delayed diagnosis and increased morbidity.

Intraoperative fluorescence angiography has recently been introduced as a means of real-time assessment of anastomotic perfusion with preliminary evidence suggesting that it may reduce the rate of AL. In addition, concepts are emerging about the role of the rectal mucosal microbiome in AL and the possible role of new prophylactic therapies.

In January 2016 a meeting of expert colorectal surgeons and pathologists was held in London, UK to identify the ongoing controversies surrounding AL in colorectal surgery. The outcome of the meeting is presented in the form of research challenges that need to be addressed.

Introduction

Anastomotic leakage (AL) is one of the most feared complications of colorectal surgery. Despite advances in anastomotic technique, postoperative monitoring and diagnostics, its incidence and consequences have not appreciably changed during the last fifty years. AL can have profound consequences for patients through mortality and short and long-term morbidity. It also has an effect on healthcare providers, through the immediate cost of management and the use of resources for remedial care. AL is considered by many surgeons to be the greatest single challenge in visceral surgery.

In January 2016, a meeting of expert colorectal surgeons and pathologists was held in London, UK to identify the challenges in preventing, diagnosing, and treating AL in light of recent innovations in this field. The meeting was sponsored by educational grants from Novadaq Technologies Inc. (Ontario, Canada) and Elemental Healthcare (Berkshire, UK). Participants were asked to give a short presentation on a specific topic related to AL, followed by an invited commentary from a second faculty member, as the basis for wider discussion. Each participant submitted a written narrative of their presentation. This manuscript, which each participant was invited to review and edit, reports the outcome of the meeting and the identified areas for further research.

Definition and Incidence

Previous research into AL has been hampered by the lack of a widely accepted definition, with twenty nine separate definitions reported from 1993 to 1999 and 59 from 2000 to 2009¹. An attempt to address this was made in 2010 by the International Study Group of Rectal Cancer (ISREC), who defined AL as ‘a communication between the intra- and extra-luminal compartments owing to a defect of the integrity of the intestinal wall at the anastomotic site’

². In addition, a pelvic abscess adjacent to an anastomosis, even if no communication with the bowel lumen could be demonstrated, was considered to have originated from a leak. Recommendation was made for a simple grading system based on clinical management as follows: grade A, resulting in no change in management, grade B, requiring active therapeutic intervention and grade C, requiring re-laparotomy or laparoscopy. Subsequent validation in patients undergoing sphincter preserving surgery of the rectum demonstrated an overall leakage rate of 7.5% with a severity of 16% grade A, 23% grade B and 61% grade C ³. Despite attempts at standardisation, AL continues to be variably reported, with an incidence ranging widely from 5% to 19%, depending on the site and type of anastomosis and the cohort under investigation ⁴. To complicate the situation further, many studies fail to take into account subclinical or radiological leakage, despite the recognised association with poor bowel function and anastomotic stricture formation ⁵.

Research challenge:

- *Integration of standard validated definition and reporting of AL into clinical trials.*

Risk factors for anastomotic leakage

Much has been written about the risk factors for AL in an attempt to guide surgeons as to when anastomosis is appropriate. In an attempt to quantitate these following left-sided colorectal surgery a Colon Leakage Score (CLS) was developed and shown to have good predictive value ⁶. The patient-related factors cited as increasing AL include male gender, smoking, obesity, alcohol abuse, pre-operative steroid and non-steroidal anti-inflammatory drugs, longer operation duration, pre-operative transfusion, contamination of the operative field and the urgency of the operation ⁴. A recent meta-analysis of 23 studies and 110,272

patients found that AL was significantly associated with low rectal anastomosis (odds ratio (OR) 3.26, 95% CI 2.31, 4.62), male gender (OR 1.48, 95% CI 1.37, 1.60), and pre-operative radiotherapy (RT) (OR 1.65, 95% CI 1.06, 2.56), although the quality of evidence was regarded as moderate or low ⁷.

The impact of RT on AL rates is a contentious issue particularly as the common practice of creating a defunctioning stoma in RT patients can introduce bias into studies ⁸. The UK Medical Research Council (MRC) CR07 randomised controlled trial (RCT) reported no difference in AL rates between patients having pre-operative RT or selective post-operative RT in a multicentre, multinational randomised analysis of 1,350 rectal cancer patients ⁹. In an interim analysis of the Stockholm III trial in rectal cancer comparing short course RT, short course RT with delay and long-course RT, the effect of radiotherapy on AL appeared to be related to the timing of surgery, with a significant increase if it was performed between 11 and 17 days following the start of RT ¹⁰.

In addition to patient-related factors and non-specialist colorectal surgeons, ¹¹ low institutional cancer case load are associated with higher AL rates ¹². A meta-analysis including 13,040 rectal cancer patients demonstrated a significantly lower risk of AL in favour of high volume hospitals (OR 0.72, 95% CI 0.48, 1.09). There was no significant difference among colon cancer patients ¹².

Research challenges:

- *Further validation of the Colon Leakage Score as a universal tool for predicting AL.*
- *Improved understanding of the detrimental effects of radiotherapy on anastomotic healing*

Anastomotic construction

The role of specialisation in reducing leakage rates highlights the importance of surgical technique in anastomotic construction. One meta-analysis of reconstruction techniques after low anterior resection compared straight coloanal anastomosis, side-to-end coloanal anastomosis, transverse coloplasty and colonic J pouch and did not demonstrate superiority of any technique in terms of leakage ¹³.

Several Cochrane reviews and meta-analyses have failed to demonstrate a benefit for either hand sewn or stapled anastomosis ¹⁴⁻¹⁷. Stapled anastomosis is increasingly being used because of technical ease, reproducibility and speed. In the era of increasing laparoscopic surgery, stapled low rectal anastomosis presents a particular problem due to the confines of the bony pelvis. Numerous studies suggest that the number of stapler firings used to transect the distal rectum in double-stapled anastomosis increases the rate of AL ¹⁸⁻²³, whilst others have failed to show this ^{24, 25}. There does not appear to be any benefit of single over double-stapled low rectal anastomosis ²⁶.

Transanal total mesorectal excision (TaTME) has been suggested to result in lower rates of AL due to the avoidance of cross stapling ^{27, 28}. More data from large international cohort studies and randomised trials are awaited to confirm this. Other techniques for anastomosis have been tried, including anastomotic compression. First described by Denans almost two centuries ago ²⁹, the AKA-2 device, developed in the 1980s, comprised two plastic rings applied with a transanal applicator. Compression of the inverted bowel edges produces simultaneous necrosis and healing. After four to six days the rings separate from the anastomosis ³⁰. The AKA-2 was superseded by the biofragmentable anastomotic ring (BAR) made of absorbable polyglycolic acid and barium sulphate ³¹. When the device is closed a

small gap remains between the rings to limit tissue necrosis. The ring fragments are passed with the stool tow to three weeks postoperatively. The limitations of the device include retained foreign material within the tissue, luminal narrowing, necrosis at the anastomotic site and problems with introducing the deployment device ³². Most recently the nickel-titanium (Nitinol; NiTi) ColonRing device has been introduced ³² with compression rings that possess shape memory and elasticity to provide more uniform compression of the tissue. Despite a large multicentre, multinational study involving 1,180 patients ³³ and a prospective multicentre study reporting leakage rates of just over 3% ³⁴ the device has not gained widespread application.

Research challenges:

- *Identification of new materials and methods for performing anastomosis with minimal collateral tissue trauma.*
- *Development of devices that accommodate the confines of the pelvis and facilitate low rectal anastomosis*

Anastomotic reinforcement

Several innovations have been tried to reinforce an anastomosis and so reduce the occurrence of leakage. These include the use of tissue adhesives, bio-absorbable materials and intraluminal stents. None has shown clinical efficacy or entered into routine practice.

Notably, two randomised comparisons of Seamguard[®] (W.L. Gore & Associates Inc., Flagstaff, Arizona, USA) against standard circular stapling for rectal anastomosis failed to show any difference in the AL rate ^{35,36}. The most recent technology, LifeSeal[™] (LifeBond, Caesarea Industrial Park, Israel), uses enzyme-based catalytic breakdown of a gelatin-based

product to seal an anastomosis; the results of early phase clinical trials are awaited. Recent studies in rat models of AL have demonstrated that cyanoacrylate-based adhesives show the greatest mechanical strength and lowest rate of ileus, bowel wall destruction and immunohistopathological tissue reaction compared with other adhesive materials^{37, 38}.

In a related strategy, omental wrap³⁹ and mesenteric flap formation⁴⁰ have been used to seal anastomoses and provide a cellular scaffold for healing. In a meta-analysis of three studies including 943 patients, the use of an omental wrap resulted in a significant three-fold reduction in the rate of clinical AL but no difference in the reoperation rate or mortality³⁹.

Research challenge:

- *Improved methods to seal an anastomosis and support tissue healing*

Defunctioning stoma

A defunctioning stoma is used to protect a rectal anastomoses by diverting the faecal stream. Whether it prevents AL or merely reduces the consequence of leakage is debated. There is no agreement whether a defunctioning stoma should be constructed for all rectal anastomoses or just for those that are low. A defunctioning ileostomy can cause morbidity, most notably dehydration and sodium depletion owing to excessive output. Reversal of the stoma is associated with a complication in some 40% of patients⁴¹. Many patients will be left with a stoma for several months due to the low clinical priority for reversal⁴² and 20% of patients are left with a permanent stoma owing to postoperative complications of anterior resection⁴³.

Several studies have suggested that a defunctioning stoma decreases the incidence of clinical leakage of a colorectal anastomosis^{44, 45}, others on the other hand have reported no difference in leakage but a reduced incidence of reoperation has been recorded⁴⁶. A recent propensity-matched scoring analysis of 936 patients who underwent low anterior resection confirms that defunctioning ileostomy does not influence the rate of clinical AL but does mitigate against the consequences, reducing the need for urgent reoperation⁴⁷.

The technique of pre-stage ileostomy, also known as ‘ghost’ ileostomy, has been described whereby a terminal ileal loop is exteriorized through the abdominal wall but is left unopened⁴⁸. Should a complication of the anastomosis develop it can be rapidly transformed into a definitive ileostomy under local anaesthesia. If this is not needed, the morbidity of an ileostomy is avoided. Although the ghost ileostomy appears to be feasible in cohort studies⁴⁹,⁵⁰, its role is yet to be determined by randomised controlled evidence.

Research challenge:

- *Improved selection criteria for patients undergoing a defunctioning stoma*
- *Further determine the role of ‘ghost’ ileostomy in association with a high risk anastomosis*

Intraoperative prediction of anastomotic leakage

The air leak test is a well established technique to check intraoperatively the anastomotic integrity immediately after completion of the anastomosis.. It is easy, cheap and quick to perform with very little risk. The evidence suggests that the air test is effective in predicting satisfactory healing when it demonstrates an intact anastomosis^{51, 52}. Thus a randomised

study by Beard *et al.*⁵¹ reported more clinical and radiological leaks in patients who did not undergo an air test than those who did (clinical 14% vs. 4%, radiological 29% vs. 11%). In a retrospective cohort study, Ricciardi *et al.*⁵² analysed 998 consecutive left sided anastomoses performed without diversion. The clinical leak rate of 8% in patients with a positive air test was significantly higher than the 4% in patients with a negative result, with the leak rate in patients with untested anastomoses being similar (8%) to those with a positive test. In the event of a positive air test, suture repair alone may be associated with a high AL rate and the evidence suggests that in this situation the anastomosis should be revised and/or defunctioned.⁵²⁻⁵⁴

Intraoperative endoscopy allows assessment of anastomotic integrity, bleeding from the staple line and the detection of any additional pathology. [52]⁵⁵. Although intra-operative endoscopy has been shown to be a safe and reliable technique, studies comparing its routine with selective use have been underpowered^{55,56}. Without a large randomised control trial any benefit of routine intraoperative endoscopy in preventing AL cannot be determined.

Research challenge:

- *A randomised evaluation of intraoperative endoscopy to assess anastomotic integrity.*

Anastomotic healing

Understanding the pathophysiology of anastomotic healing is crucial to the development of new treatment and preventative strategies. Owing to the complex biological processes involved, it is difficult to mimic anastomotic healing *in vitro* and a recent systematic review showed that animal research on AL is of poor quality⁵⁷. Although many components of the

healing process, such as the initial inflammatory response, organisation, collagen deposition and wound remodelling, are common to all tissues and have been studied extensively in skin, certain aspects of anastomotic healing differ considerably⁵⁸. Caution is therefore advised in translating cutaneous models of wound healing to healing in the gastro-intestinal (GI) tract.

Collagen in the GI tract is produced both by fibroblasts and smooth muscle cells compared with fibroblasts alone in the skin. Despite increased shear stress and bacterial load, wound healing is far more rapid in the GI tract⁵⁸. The capacity of collagen synthesis in anastomotic tissue⁵⁹ is significantly impaired in sepsis and an anastomosis should be avoided in such circumstances. Furthermore, collagenase, which is less important in cutaneous healing, plays a critical role. Collagenase is upregulated in the colonic mucosa following anastomosis, contributing to lysis of collagen and decreased anastomotic strength⁶⁰. It has been suggested that some patients are at increased risk of AL due to pre-existing changes in the extracellular matrix, such as a lower collagen type I/III ratio and higher expression of several collagenase subtypes in both mucosa and submucosa⁶¹.

Research challenge:

- *Improved understanding of the molecular and biochemical pathways controlling the early balance of collagen synthesis and breakdown in anastomotic healing*

Optimising anastomotic vascularity

Optimal blood supply to the anastomosis is critical for healing. At operation, perfusion is assessed clinically by inspecting the colour of the bowel and bleeding from its cut ends and the marginal vessels. Several attempts have been made to apply more sophisticated means of

assessing anastomotic perfusion. Such methods include analysis of tissue oxygenation using light spectroscopy⁶² and modified pulse oximetry^{63, 64}. A fall in tissue oxygen tension appears to be associated with subsequent AL. Interestingly, in patients who do not suffer AL there is a reactive rise in the tissue oxygen tension in the proximal part of the anastomosis, suggesting a protective compensatory mechanism at the tissue level⁶².

In patients undergoing chemoradiotherapy for rectal cancer, computed tomographic (CT) perfusion scanning was used to assess parameters of microvascular perfusion, including blood flow and blood volume⁶⁵. Both parameters were increased in the cancers compared with normal rectal wall and showed a significant reduction in response to chemoradiotherapy. Similar techniques may be applicable to assessing the risk of AL particularly after radiotherapy.

The development of fluorescence perfusion angiography has enabled the surgeon to 'visualise' tissue perfusion in real time. This allows the surgeon to assess bowel perfusion and to select an appropriate segment of bowel for anastomosis and then to check perfusion endoscopically on completion of the anastomosis⁶⁶. Typically, a fluorophore is injected intravenously and visualized with a near-infrared (NIR) camera. NIR imaging has the advantage over other wavelengths of electromagnetic radiation in providing deep tissue penetration without causing thermal damage⁶⁷. The most commonly used fluorophore is indocyanine green (ICG), which absorbs light at 790-810nm and re-emits fluorescence at 835nm⁶⁷. ICG is safe for human use and has US Food and Drug Administration (FDA) approval for a variety of imaging applications, including assessment of hepatic function, measurement of cardiac output, and ophthalmic angiography⁶⁸. When administered intravenously it binds rapidly to plasma proteins and is limited to the intravascular

compartment. It has a plasma elimination half-life of three to five minutes and is metabolized in the liver with a half-life of 15 to 20 minutes, making it suitable for multiple evaluations^{69, 70}.

Intra-operative NIR-ICG for assessment of anastomotic perfusion has been shown to be feasible and safe in several non-randomised studies using laparoscopic (PINPOINT™ (Novadaq, Mississauga, Ontario, Canada)^{69, 71}, NIR/ICG system (Karl Storz⁷², Tuttlingen, Germany)) and robotic (Firefly™ (Intuitive Surgical Inc., Sunnyvale, California, USA)) systems⁷³. In white light mode these systems act as a conventional laparoscope and in NIR mode the ICG is visualized as a white fluorescence on a black background. The PINPOINT™ system provides a dual display mode, where green ICG fluorescence is superimposed on the white light image to provide an augmented view of tissue perfusion. All commercially available NIR imaging systems lack the capability to quantify tissue perfusion, which remains subjective with no clear cut-off to guide the surgeon on the adequacy or inadequacy of anastomotic perfusion⁶⁷.

The largest non-randomised multicentre trial to use NIR fluorescence angiography with the PINPOINT system to assess anastomotic viability during laparoscopic left colectomy is the Perfusion Assessment in Laparoscopic Left Anterior Resection (PILLAR II) trial⁷¹. This feasibility study in 147 patients conducted in the United States achieved successful imaging in 98.6% of cases and resulted in a change of surgical plan in 7.9% of the patients. There was a low clinical leak rate of 1.4% at 30-days with no complications attributable to the use of ICG or the device. A European phase II study of the role of NIR fluorescence angiography in assessing anastomotic perfusion is ongoing (clinical trials identifier NCT02459405)⁷⁴.

A RCT evaluating ICG-NIR in rectal surgery, the PILLAR III trial, is due for completion in 2017 (clinical trials identifier NCT02205307) ⁷⁵. Patients planned for open or minimally invasive low anterior resection for rectal or rectosigmoid cancer with an anastomosis at 10cm or less from the anal verge, are eligible. They are stratified according to neo-adjuvant therapy and randomised to receive intra-operative fluorescent angiography with the PINPOINT[®] or SPY[®] Elite system or white light laparoscopy. The primary endpoint is the anastomotic leakage rate.

Research challenges:

- *Validation of NIR-ICG efficacy in reducing AL*
- *Methods to quantify NIR-ICG perfusion*
- *Multi-spectral laparoscopic systems for detection of different fluorophores*
- *Novel pharmacological methods to enhance anastomotic perfusion*

Anastomotic leakage as an infective complication

There has been much interest in a return to the concept of anastomotic leakage as an infective complication. Using a rat model, Shogan *et al* have shown that anastomotic injury results in a change in anastomotic tissue-associated microbiota with a notable 500-fold and 200-fold increase in the relative abundance of *Enterococcus* and *Escherichia/Shigella* species ⁷⁶. Importantly, this difference was only apparent in tissue from the anastomosis and not in luminal faecal samples. AL was associated with increased bacterial virulence-associated pathways, including production of matrix-degrading enzymes and cytotoxic necrotizing factors. Work by the same group, again in a rat model, showed that *Enterococcus faecalis* contributed to AL by upregulation of collagenase activity and activation of tissue matrix metalloproteinase-9 (MMP-9) and that AL was prevented by administration of an antibiotic

enema or MMP-9 inhibitor ⁷⁷. Furthermore, in a small cohort of 11 patients undergoing colonic surgery, *E. faecalis* and other bacteria with collagen-degrading and MMP-9-activating ability could be isolated from the anastomotic site and were unaffected by the use of standard intravenous prophylactic antibiotics.

The concept of upregulation of bacterial virulence pathways with increased collagenase production leading to AL is supported by work from the 1980s conducted by Young and Wheeler ^{78, 79}. Using a collagenase inhibitor (Aprotinin) they were able to demonstrate an increase in anastomotic burst pressure and breaking strength in a rat model ⁷⁸ and a significant reduction of radiological and clinical AL in a RCT involving 100 patients ⁷⁹.

Another interesting observation, with relevance to rectal cancer surgery, is the change in composition and virulence of the rectal flora following radiotherapy ⁸⁰. The adverse influence of radiotherapy on AL is usually attributed to tissue inflammation and microvascular injury, but it is possible that radiotherapy-induced changes in the rectal flora result in a pro-AL microenvironment. This is supported by the work of Olivas *et al* (2012) who showed in a model of low anterior resection that pre-operative radiation and intestinal inoculation of *Pseudomonas aeruginosa* (a collagenase producing bacterium) resulted in high rates of AL, whereas radiation alone or *P. aeruginosa* alone did not cause leakage ⁸¹.

Research challenges

- *To establish cause and effect for the role of the microbiome in AL*
- *Exploit protease inhibition as a mechanism to reduce AL*
- *Further investigation of the change in microbiome induced by radiation, its role in AL, and strategies for mitigation.*

The role of mechanical bowel preparation

The role of mechanical bowel preparation (MBP) prior to surgery is an area of contention, with opinions varying either side of the Atlantic. The disadvantages of MBP are the unwanted complications, including hypovolaemia, metabolic disturbance and poor patient compliance⁸²⁻⁸⁵. The advantages include the ability to perform intraoperative endoscopy and to avoid a faecally loaded colon. Evidence against the use of routine MBP in colonic surgery includes a meta-analysis performed in 2009 and a subsequent Cochrane review^{86, 87}.

The evidence for MBP in rectal surgery is less clear-cut. Randomised evidence in rectal cancer is confined to one trial, GRECCAR III, which randomised patients to MBP (oral laxative and enema) or no MBP⁸⁴. Although no difference in the rates of AL and major morbidity were demonstrated, patients with no MBP had a higher 30-day morbidity and more infectious complications. Other studies have confirmed the lack of association between MBP and AL in rectal surgery^{88, 89} and suggested that a rectal enema alone may be as effective⁹⁰. This is the focus of an ongoing Italian RCT in which patients are randomised to either full MBP or rectal enema alone (clinical trials identifier NCT00940030)⁷⁴.

Research challenge:

- *Optimal method of bowel cleansing prior to rectal surgery*
- *Clarification on the role of MBP when an upstream stoma is to be fashioned*

Pre-operative antibiotics and selective gut decontamination

The pre-operative administration of non-absorbable oral antibiotic preparations (OAP) has been demonstrated to reduce the risk of AL following elective GI surgery (OR 0.42, 95% CI 0.24 to 0.73)⁹¹. The co-administration of MBP with OAP may reduce the faecal load allowing increased delivery of OAP to the colonic mucosa^{92, 93}. Several recent large retrospective studies using the National Surgical Quality Improvement Program (NSQIP) database suggest that OAP in combination with MBP is effective in reducing AL⁹³⁻⁹⁵. There are, however, methodological limitations with these studies, including differences in the study populations, lack of data on systemic antibiotic usage and lack of detail regarding type and dose of MBP and OAP. Further evidence is needed and might be forthcoming from the SELECT study (clinical trials identifier NCT01740947), which is investigating whether oral non-absorbable antibiotics reduce clinical AL, and is due to complete in 2018⁹⁶.

Research challenge:

- *The optimal administration and antibiotic combination for reducing AL and surgical site infections*

Early diagnosis

Early diagnosis of anastomotic leakage allows timely remedial intervention and reduces morbidity and mortality⁹⁷. The difficulty is that AL presents in a variety of ways, ranging from rapid fulminant sepsis to a more insidious onset with failure to progress in the postoperative period. The main radiological modalities used to diagnose AL are water-soluble contrast enema (WSCE) and CT, but data on their accuracy are limited by the timing of the study and the expertise of the radiologist⁹⁸. WSCE is safe despite fears concerning

anastomotic disruption⁹⁹, but its sensitivity and specificity can be poor with reported figures of 52% and 87% respectively in left-sided anastomoses¹⁰⁰. In comparison, the sensitivity and specificity of CT scanning has been reported to be 57% and 100% after total mesorectal excision for rectal cancer¹⁰¹. It is possible that a combination of the techniques, using CT with rectal contrast, improves diagnostic accuracy¹⁰².

The use of biosensors to sample the anastomotic environment via an intraluminal or microdialysis catheter may allow early detection of anastomotic disruption⁵⁵. This technique allows rapid analysis at multiple time points. Studies have reported a lower pH¹⁰³ and concentration of ischaemia-related metabolites¹⁰⁴ and increased levels of lysozyme¹⁰⁵, bacterial cell wall lipopolysaccharide¹⁰⁶ and cytokines¹⁰⁷ in patients who develop AL. These findings suggest that bacterial translocation has occurred within the first three to five days of the anastomosis being constructed, even if the frank leakage of intraluminal contents has not occurred by that time.

Changes in electrical resistance have also been associated with anastomotic disruption¹⁰⁸. The use of serum C-reactive protein (CRP) as a marker of AL is probably limited by its poor specificity, but it has been suggested that a CRP above 150mmol/l on postoperative days 3 to 5 may be predictive and a trigger for further investigation⁴.

Research challenges:

- *The optimal radiological techniques to detect AL*
- *Identification of the preferred biomarkers for detecting early AL*
- *Development of real-time, bed-side methods for detecting early AL*

Management strategies for anastomotic leakage

Many factors have to be considered when deciding the best treatment for a patient with AL. These include age, comorbidity, level of the anastomosis, the interval from the primary operation and the degree of anastomotic dehiscence ¹⁰⁹. The treatment options include one or more of antibiotic therapy, transanal drainage, percutaneous drainage, laparoscopy/laparotomy with anastomotic repair and defunctioning stoma, and take-down of the anastomosis with end stoma. A survey of 350 members of the Dutch Society of Gastrointestinal Surgery suggested heterogeneity in the strategies employed to manage leakage with a preference to try and preserve a left sided anastomosis in fitter and younger patients ¹¹⁰.

In an analysis of the American College of Surgeons (ACS) NSQIP 2012 database, 56% of patients with AL required re-operation, 25% received a non-surgical intervention and 19% were treated medically ¹¹¹. The need for re-operation was found to be greater in patients with anastomotic leakage from a colo-colonic anastomosis than with an ileo-colonic or colo-rectal anastomosis, and was reduced in patients with a stoma. In patients with a high-risk profile, a repair or construction of new anastomosis should probably not be performed without a protective stoma ¹¹². McDermott et al proposed an AL severity score to aid clinical decision-making regarding the escalation of care, use of inotropes and intervention ⁴. This score ranges from a patient deviating from his or her expected course with biochemical abnormalities (grade 1), where the recommendation is close observation to a patient in septic shock (grade 5) where intensive care and emergency laparotomy are required.

AL following rectal surgery, particularly in the presence of a diverting stoma, may present as a pre-sacral abscess that can leave a chronic sinus after it has drained. Treatment options include a watch and wait policy, take down of the anastomosis and creation of an end colostomy or laying open of the sinus into the neorectum. Newer techniques include endoluminal endosponge vacuum-assisted systems, which provide continuous drainage and encourage by suction the bowel wall of the neorectum to obliterate the cavity¹¹³. This should, however, be undertaken early whilst the tissues are still flexible and can respond to the negative pressure. Vacuum—assisted treatment is effective with and without a diverting stoma^{113, 114}. The use of sutures or endoscopic clips to close the residual defect following endosponge therapy may reduce the time to healing¹¹⁵.

Research challenges

- *Validation of endoluminal therapies for anastomotic leakage*

Summary and future perspectives

AL is a substantial problem for clinicians, patients and healthcare providers. Little progress has been made over the last ten years in our understanding or management of the condition.

A concerted effort is required by surgical researchers, working in collaboration with industry, basic scientists and healthcare partners. Several new techniques have emerged that hold promise in reducing AL although their exact role is yet to be established by prospective randomised evidence. The emerging concept of AL as an infective complication warrants further investigation and if it is shown to be valid it may hold the key to future prophylactic therapies.

The aim of this manuscript has been to present the current thinking amongst leading colorectal surgeons and pathologists on the most pressing areas for further research. It is hoped that this manuscript will stimulate the research that is needed to make an impact on a clinical problem that has defied the efforts of previous generations.

Acknowledgements

The following individuals also participated in the expert meeting on colorectal anastomotic leak: B. George (Oxford University Hospitals, UK), M. Chand (University College Hospital, UK), M. Reif (Novadaq Technologies Inc., Canada) and T. Yeung (University of Oxford, UK).

References

1. Bruce J, Krukowski ZH, Al-Khairy G, Russell EM, Park KG. Systematic review of the definition and measurement of anastomotic leak after gastrointestinal surgery. *Br J Surg*. 2001;88(9):1157-68.
2. Rahbari NN, Weitz J, Hohenberger W, Heald RJ, Moran B, Ulrich A, et al. Definition and grading of anastomotic leakage following anterior resection of the rectum: a proposal by the International Study Group of Rectal Cancer. *Surgery*. 2010;147(3):339-51.
3. Kulu Y, Ulrich A, Bruckner T, Contin P, Welsch T, Rahbari NN, et al. Validation of the International Study Group of Rectal Cancer definition and severity grading of anastomotic leakage. *Surgery*. 2013;153(6):753-61.
4. McDermott FD, Heeney A, Kelly ME, Steele RJ, Carlson GL, Winter DC. Systematic review of pre-operative, intraoperative and postoperative risk factors for colorectal anastomotic leaks. *British Journal of Surgery*. 2015;102(5):462-79.
5. Lim M, Akhtar S, Sasapu K, Harris K, Burke D, Sagar P, et al. Clinical and subclinical leaks after low colorectal anastomosis: a clinical and radiologic study. *Dis Colon Rectum*. 2006;49(10):1611-9.
6. Dekker JW, Liefers GJ, de Mol van Otterloo JC, Putter H, Tollenaar RA. Predicting the risk of anastomotic leakage in left-sided colorectal surgery using a colon leakage score. *Journal of Surgical Research*. 2011;166(1):e27-34.
7. Pommergaard HC, Gessler B, Burcharth J, Angenete E, Haglund E, Rosenberg J. Pre-operative risk factors for anastomotic leakage after resection for colorectal cancer: a systematic review and meta-analysis. *Colorectal Disease*. 2014;16(9):662-71.
8. Marijnen CAM, Kapiteijn E, van de Velde CJH, Martijn H, Steup WH, Wiggers T, et al. Acute Side Effects and Complications After Short-Term Pre-operative Radiotherapy Combined With Total Mesorectal Excision in Primary Rectal Cancer: Report of a Multicenter Randomized Trial. *Journal of Clinical Oncology*. 2002;20(3):817-25.

9. Sebag-Montefiore D, Stephens RJ, Steele R, Monson J, Grieve R, Khanna S, et al. Pre-operative radiotherapy versus selective post-operative chemoradiotherapy in patients with rectal cancer (MRC CR07 and NCIC-CTG C016): a multicentre, randomised trial. *Lancet*. 2009;373(9666):811-20.
10. Pettersson D, Cedermark B, Holm T, Radu C, Pahlman L, Glimelius B, et al. Interim analysis of the Stockholm III trial of pre-operative radiotherapy regimens for rectal cancer. *British Journal of Surgery*. 2010;97(4):580-7.
11. Biondo S, Kreisler E, Millan M, Fraccalvieri D, Golda T, Frago R, et al. Impact of surgical specialization on emergency colorectal surgery outcomes. *Archives of Surgery*. 2010;145(1):79-86.
12. Archampong D, Borowski D, Wille-Jorgensen P, Iversen LH. Workload and surgeon's specialty for outcome after colorectal cancer surgery. *Cochrane Database Syst Rev*. 2012;3:Cd005391.
13. Hüttner FJ, Tenckhoff S, Jensen K, Uhlmann L, Kulu Y, Büchler MW, et al. Meta-analysis of reconstruction techniques after low anterior resection for rectal cancer. *British Journal of Surgery*. 2015;102(7):735-45.
14. Choy PY, Bissett IP, Docherty JG, Parry BR, Merrie AE. Stapled versus handsewn methods for ileocolic anastomoses. *Cochrane Database Syst Rev*. 2007(3):Cd004320.
15. Lustosa SA, Matos D, Atallah AN, Castro AA. Stapled versus handsewn methods for colorectal anastomosis surgery. *Cochrane Database Syst Rev*. 2001(3):Cd003144.
16. MacRae HM, McLeod RS. Handsewn vs. stapled anastomoses in colon and rectal surgery: a meta-analysis. *Dis Colon Rectum*. 1998;41(2):180-9.
17. Sliker JC, Daams F, Mulder IM, Jeekel J, Lange JF. Systematic review of the technique of colorectal anastomosis. *JAMA Surgery*. 2013;148(2):190-201.
18. Ito M, Sugito M, Kobayashi A, Nishizawa Y, Tsunoda Y, Saito N. Relationship between multiple numbers of stapler firings during rectal division and anastomotic leakage after laparoscopic rectal resection. *Int J Colorectal Dis*. 2008;23(7):703-7.
19. Kawada K, Hasegawa S, Hida K, Hirai K, Okoshi K, Nomura A, et al. Risk factors for anastomotic leakage after laparoscopic low anterior resection with DST anastomosis. *Surgical Endoscopy*. 2014;28(10):2988-95.
20. Kim CW, Baek SJ, Hur H, Min BS, Baik SH, Kim NK. Anastomotic Leakage After Low Anterior Resection for Rectal Cancer Is Different Between Minimally Invasive Surgery and Open Surgery. *Ann Surg*. 2016;263(1):130-7.
21. Kim JS, Cho SY, Min BS, Kim NK. Risk factors for anastomotic leakage after laparoscopic intracorporeal colorectal anastomosis with a double stapling technique. *Journal of the American College of Surgeons*. 2009;209(6):694-701.
22. Park JS, Choi G-S, Kim SH, Kim HR, Kim NK, Lee KY, et al. Multicenter analysis of risk factors for anastomotic leakage after laparoscopic rectal cancer excision: the Korean laparoscopic colorectal surgery study group. *Ann Surg*. 2013;257(4):665-71.
23. Katsuno H, Shiomi A, Ito M, Koide Y, Maeda K, Yatsuoka T, et al. Comparison of symptomatic anastomotic leakage following laparoscopic and open low anterior resection for rectal cancer: a propensity score matching analysis of 1014 consecutive patients. *Surgical Endoscopy*. 2016;30(7):2848-56.
24. Akiyoshi T, Ueno M, Fukunaga Y, Nagayama S, Fujimoto Y, Konishi T, et al. Incidence of and risk factors for anastomotic leakage after laparoscopic anterior resection with intracorporeal rectal transection and double-stapling technique anastomosis for rectal cancer. *American Journal of Surgery*. 2011;202(3):259-64.
25. Yamamoto S, Fujita S, Akasu T, Inada R, Moriya Y. Risk factors for anastomotic leakage after laparoscopic surgery for rectal cancer using a stapling technique. *Surgical Laparoscopy Endoscopy & Percutaneous Techniques*. 2012;22(3):239-43.

26. Kim HJ, Choi GS, Park JS, Park SY. Comparison of intracorporeal single-stapled and double-stapled anastomosis in laparoscopic low anterior resection for rectal cancer: a case-control study. *Int J Colorectal Dis.* 2013;28(1):149-56.
27. Penna M, Knol J, Tuynman J, Tekkis P, Mortensen N, Hompes R. Four anastomotic techniques following transanal total mesorectal excision (TaTME). *Tech Coloproctol.* 2016;20(3):185-91.
28. Simillis C, Hompes R, Penna M, Rasheed S, Tekkis PP. A systematic review of transanal total mesorectal excision: is this the future of rectal cancer surgery? *Colorectal Disease.* 2016;18(1):19-36.
29. Denans F. Nouveau procédé pour la guérison des plaies des intestins: recueil de la société Royale de Médecine de Marseille. Imprimerie d'Archard; Marseille, France 1827.
30. Kaidar-Person O, Rosenthal RJ, Wexner SD, Szomstein S, Person B. Compression anastomosis: history and clinical considerations. *Am J Surg.* 2008;195(6):818-26.
31. Kim S-H, Choi H-J, Park K-J, Kim J-M, Kim K-H, Kim M-C, et al. Sutureless Intestinal Anastomosis With the Biofragmentable Anastomosis Ring: Experience of 632 Anastomoses in a Single Institute. *Diseases of the Colon & Rectum.* 2005;48(11):2127-32.
32. Buchberg BS, Masoomi H, Bergman H, Mills SD, Stamos MJ. The Use of a Compression Device as an Alternative to Hand-Sewn and Stapled Colorectal Anastomoses: Is Three a Crowd? *Journal of Gastrointestinal Surgery.* 2011;15(2):304-10.
33. Masoomi H, Luo R, Mills S, Carmichael JC, Senagore AJ, Stamos MJ. Compression anastomosis ring device in colorectal anastomosis: a review of 1,180 patients. *The American Journal of Surgery.* 2013;205(4):447-51.
34. D'Hoore A, Albert MR, Cohen SM, Herbst F, Matter I, Van Der Speeten K, et al. COMPRES: a prospective postmarketing evaluation of the compression anastomosis ring CAR 27 /ColonRing. *Colorectal Dis.* 2015;17(6):522-9.
35. Senagore A, Lane FR, Lee E, Wexner S, Dujovny N, Sklow B, et al. Bioabsorbable staple line reinforcement in restorative proctectomy and anterior resection: a randomized study. *Diseases of the Colon & Rectum.* 2014;57(3):324-30.
36. Placer C, Enriquez-Navascues JM, Elorza G, Timoteo A, Mugica JA, Borda N, et al. Preventing complications in colorectal anastomosis: results of a randomized controlled trial using bioabsorbable staple line reinforcement for circular stapler. *Dis Colon Rectum.* 2014;57(10):1195-201.
37. Vakalopoulos KA, Wu Z, Kroese L, Kleinrensink GJ, Jeekel J, Vendamme R, et al. Mechanical strength and rheological properties of tissue adhesives with regard to colorectal anastomosis: an ex vivo study. *Ann Surg.* 2015;261(2):323-31.
38. Vakalopoulos KA, Wu Z, Kroese LF, van der Horst PH, Lam KH, Dodou D, et al. Clinical, mechanical, and immunohistopathological effects of tissue adhesives on the colon: An in-vivo study. *Journal of Biomedical Materials Research Part B: Applied Biomaterials.* 2016:n/a-n/a.
39. Hao XY, Guo TK, Ma B, Tian JH, Li HL. Omentoplasty in the prevention of anastomotic leakage after colorectal resection: a meta-analysis. *In J Colorect Dis.* 2008;23:1159 - 65.
40. Mohan HM, Winter DC. Autobuttrussing of colorectal anastomoses using a mesenteric flap. *Updates in Surgery.* 2013;65(4):333-5.
41. Akesson O, Syk I, Lindmark G, Buchwald P. Morbidity related to defunctioning loop ileostomy in low anterior resection. *Int J Colorectal Dis.* 2012;27(12):1619-23.
42. Floodeen H, Lindgren R, Matthiessen P. When are defunctioning stomas in rectal cancer surgery really reversed? Results from a population-based single center experience. *Scandinavian Journal of Surgery: SJS.* 2013;102(4):246-50.

43. Lindgren R, Hallbook O, Rutegard J, Sjodahl R, Matthiessen P. What is the risk for a permanent stoma after low anterior resection of the rectum for cancer? A six-year follow-up of a multicenter trial. *Diseases of the Colon & Rectum*. 2011;54(1):41-7.
44. Chude GG, Rayate NV, Patris V, Koshariya M, Jagad R, Kawamoto J, et al. Defunctioning loop ileostomy with low anterior resection for distal rectal cancer: should we make an ileostomy as a routine procedure? A prospective randomized study. *Hepato-Gastroenterology*. 2008;55(86-87):1562-7.
45. Tan WS, Tang CL, Shi L, Eu KW. Meta-analysis of defunctioning stomas in low anterior resection for rectal cancer. *British Journal of Surgery*. 2009;96(5):462-72.
46. Matthiessen P, Hallbook O, Rutegard J, Simert G, Sjodahl R. Defunctioning stoma reduces symptomatic anastomotic leakage after low anterior resection of the rectum for cancer: a randomized multicenter trial. *Annals of Surgery*. 2007;246(2):207-14.
47. Shiomi A, Ito M, Maeda K, Kinugasa Y, Ota M, Yamaue H, et al. Effects of a diverting stoma on symptomatic anastomotic leakage after low anterior resection for rectal cancer: a propensity score matching analysis of 1,014 consecutive patients. *J Am Coll Surg*. 2015;220(2):186-94.
48. Miccini M, Amore Bonapasta S, Gregori M, Barillari P, Tocchi A. Ghost ileostomy: real and potential advantages. *Am J Surg*. 2010;200(4):e55-7.
49. Cerroni M, Cirocchi R, Morelli U, Trastulli S, Desiderio J, Mezzacapo M, et al. Ghost Ileostomy with or without abdominal parietal split. *World Journal of Surgical Oncology*. 2011;9(1):1-5.
50. Mori L, Vita M, Razzetta F, Meinero P, D'Ambrosio G. Ghost Ileostomy in Anterior Resection for Rectal Carcinoma: Is It Worthwhile? *Diseases of the Colon & Rectum*. 2013;56(1):29-34.
51. Beard JD, Nicholson ML, Sayers RD, Lloyd D, Everson NW. Intraoperative air testing of colorectal anastomoses: a prospective, randomized trial. *Br J Surg*. 1990;77(10):1095-7.
52. Ricciardi R, Roberts PL, Marcello PW, Hall JF, Read TE, Schoetz DJ. Anastomotic leak testing after colorectal resection: what are the data? *Arch Surg*. 2009;144(5):407-11; discussion 11-2.
53. Kamal T, Pai A, Velchuru VR, Zawadzki M, Park JJ, Marecik SJ, et al. Should anastomotic assessment with flexible sigmoidoscopy be routine following laparoscopic restorative left colorectal resection? *Colorectal Disease*. 2015;17(2):160-4.
54. Li VK, Wexner SD, Pulido N, Wang H, Jin HY, Weiss EG, et al. Use of routine intraoperative endoscopy in elective laparoscopic colorectal surgery: can it further avoid anastomotic failure? *Surg Endosc*. 2009;23(11):2459-65.
55. Hirst NA, Tiernan JP, Millner PA, Jayne DG. Systematic review of methods to predict and detect anastomotic leakage in colorectal surgery. *Colorectal Disease*. 2014;16(2):95-109.
56. Ishihara S, Watanabe T, Nagawa H. Intraoperative colonoscopy for stapled anastomosis in colorectal surgery. *Surg Today*. 2008;38(11):1063-5.
57. Yauw STK, Wever KE, Hoesseini A, Ritskes-Hoitinga M, van Goor H. Systematic review of experimental studies on intestinal anastomosis. *British Journal of Surgery*. 2015;102(7):726-34.
58. Thornton FJ, Barbul A. HEALING IN THE GASTROINTESTINAL TRACT. *Surgical Clinics of North America*. 1997;77(3):549-73.
59. Ahrendt GM, Tantry US, Barbul A. Intra-abdominal sepsis impairs colonic reparative collagen synthesis. *The American Journal of Surgery*. 1996;171(1):102-8.
60. Hawley PR, Faulk WP, Hunt TK, Dunphy JE. Collagenase activity in the gastrointestinal tract. *British Journal of Surgery*. 1970;57(12):896-900.

61. Stumpf M, Klinge U, Wilms A, Zabrocki R, Rosch R, Junge K, et al. Changes of the extracellular matrix as a risk factor for anastomotic leakage after large bowel surgery. *Surgery*. 2005;137(2):229-34.
62. Karliczek A, Benaron DA, Baas PC, Zeebregts CJ, Wiggers T, van Dam GM. Intraoperative assessment of microperfusion with visible light spectroscopy for prediction of anastomotic leakage in colorectal anastomoses. *Colorectal Disease*. 2010;12(10):1018-25.
63. Myers C, Mutafyan G, Petersen R, Pryor A, Reynolds J, Demaria E. Real-time probe measurement of tissue oxygenation during gastrointestinal stapling: mucosal ischemia occurs and is not influenced by staple height. *Surgical Endoscopy*. 2009;23(10):2345-50.
64. Servais EL, Rizk NP, Oliveira L, Rusch VW, Bikson M, Adusumilli PS. Real-time intraoperative detection of tissue hypoxia in gastrointestinal surgery by wireless pulse oximetry. *Surgical Endoscopy*. 2011;25(5):1383-9.
65. Bellomi M, Petralia G, Sonzogni A, Zampino MG, Rocca A. CT perfusion for the monitoring of neoadjuvant chemotherapy and radiation therapy in rectal carcinoma: initial experience. *Radiology*. 2007;244(2):486-93.
66. Jafari MD, Lee KH, Halabi WJ, Mills SD, Carmichael JC, Stamos MJ, et al. The use of indocyanine green fluorescence to assess anastomotic perfusion during robotic assisted laparoscopic rectal surgery. *Surg Endosc*. 2013;27(8):3003-8.
67. Cahill RA, Ris F, Mortensen NJ. Near-infrared laparoscopy for real-time intraoperative arterial and lymphatic perfusion imaging. *Colorectal Disease*. 2011;13:12-7.
68. Fengler J. Near-infrared fluorescence laparoscopy- technical description of PINPOINT® a novel and commercially available system. *Colorectal Disease*. 2015;17:3-6.
69. Ris F, Hompes R, Cunningham C, Lindsey I, Guy R, Jones O, et al. Near-infrared (NIR) perfusion angiography in minimally invasive colorectal surgery. *Surgical Endoscopy*. 2014;28(7):2221-6.
70. Cahill RA, NJ. M. Intraoperative augmented reality for laparoscopic colorectal surgery by intraoperative near-infrared fluorescence imaging and optical coherence tomography. *Minerva Chir*. 2011;65(4):451-62.
71. Jafari MD, Wexner SD, Martz JE, McLemore EC, Margolin DA, Sherwinter DA, et al. Perfusion assessment in laparoscopic left-sided/anterior resection (PILLAR II): a multi-institutional study. *J Am Coll Surg*. 2015;220(1):82-92.e1.
72. Boni L, David G, Mangano A, Dionigi G, Rausei S, Spampatti S, et al. Clinical applications of indocyanine green (ICG) enhanced fluorescence in laparoscopic surgery. *Surg Endosc*. 2015;29(7):2046-55.
73. Hellan M, Spinoglio G, Pigazzi A, Lagares-Garcia JA. The influence of fluorescence imaging on the location of bowel transection during robotic left-sided colorectal surgery. *Surg Endosc*. 2014;28(5):1695-702.
74. ClinicalTrials.gov. Decreasing Leak Rate in Colorectal Surgery Using Near Infra-red (NIR) Imaging 2013 [04/04/2016]. Available from: <https://clinicaltrials.gov/ct2/show/NCT02459405>
75. ClinicalTrials.gov. A Study Assessing Perfusion Outcomes With PINPOINT® Near Infrared Fluorescence Imaging in Low Anterior Resection (PILLAR III) 2015 [05/02/2016]. Available from: <https://clinicaltrials.gov/ct2/show/NCT02205307>.
76. Shogan BD SD, Chritley S, Gilbert JA, Zaborina O, Alverdy JC. Intestinal anastomotic injury alters spatially defined microbiome composition and function. *Microbiome*. 2014;2:35.
77. Shogan BD, Belogortseva N, Luong PM, Zaborin A, Lax S, Bethel C, et al. Collagen degradation and MMP9 activation by *Enterococcus faecalis* contribute to intestinal anastomotic leak. *Science Translational Medicine*. 2015;7(286):286ra68.

78. Young HL WM. Collagenase inhibition in the healing colon. *J Roy Soc Med.* 1983;76:5.
79. Young HL WM. Results of a prospective randomized double-blind trial of aprotinin in colonic surgery. *World J Surg.* 1984;8:8.
80. Brook I WR, MacVittie TJ. Effect of antimicrobial therapy on bowel flora and bacterial infection in irradiated mice. *Int J Rad Biol.* 1988;53:8.
81. Olivas AD, Shogan BD, Valuckaite V, Zaborin A, Belogortseva N, Musch M, et al. Intestinal tissues induce an SNP mutation in *Pseudomonas aeruginosa* that enhances its virulence: possible role in anastomotic leak. *PLoS ONE [Electronic Resource]*. 2012;7(8):e44326.
82. Kim YS, Hong CW, Kim BC, Han KS, Park JW, Seong Choi H, et al. Randomized clinical trial comparing reduced-volume oral picosulfate and a prepackaged low-residue diet with 4-liter PEG solution for bowel preparation. *Dis Colon Rectum.* 2014;57(4):522-8.
83. Hookey LC, Depew WT, Vanner S. The safety profile of oral sodium phosphate for colonic cleansing before colonoscopy in adults. *Gastrointestinal Endoscopy.* 2005;56(6):895-902.
84. Bretagnol F, Panis Y, Rullier E, Rouanet P, Berdah S, Dousset B, et al. Rectal cancer surgery with or without bowel preparation: The French GRECCAR III multicenter single-blinded randomized trial. *Ann Surg.* 2010;252(5):863-8.
85. Shapira Z, Feldman L, Lavy R, Weissgarten J, Haitov Z, Halevy A. Bowel preparation: comparing metabolic and electrolyte changes when using sodium phosphate/polyethylene glycol. *Int J Surg.* 2010;8(5):356-8.
86. Slim K, Vicaut E, Launay-Savary MV, Contant C, Chipponi J. Updated systematic review and meta-analysis of randomized clinical trials on the role of mechanical bowel preparation before colorectal surgery. *Ann Surg.* 2009;249(2):203-9.
87. Guenaga KF, Matos D, Wille-Jorgensen P. Mechanical bowel preparation for elective colorectal surgery. *Cochrane Database Syst Rev.* 2011(9):Cd001544.
88. van't Sant HP, Weidema WF, Hop WC, Lange JF, Contant CM. The influence of mechanical bowel preparation in elective lower colorectal surgery. *Annals of Surgery.* 2010;252(3):575-6.
89. Courtney DE, Kelly ME, Burke JP, Winter DC. Postoperative outcomes following mechanical bowel preparation before proctectomy: a meta-analysis. *Colorectal Disease.* 2015;17(10):862-9.
90. Pittet O, Nocito A, Balke H, Duvoisin C, Clavien PA, Demartines N, et al. Rectal enema is an alternative to full mechanical bowel preparation for primary rectal cancer surgery. *Colorectal Dis.* 2015;17(11):1007-10.
91. Roos D, Dijkman LM, Tijssen JG, Gouma DJ, Gerhards MF, Oudemans-van Straaten HM. Systematic review of perioperative selective decontamination of the digestive tract in elective gastrointestinal surgery. *British Journal of Surgery.* 2013;100(12):1579-88.
92. Fry DE. Colon preparation and surgical site infection. *Am J Surg.* 2011;202(2):225-32.
93. Scarborough JE, Mantyh CR, Sun Z, Migaly J. Combined Mechanical and Oral Antibiotic Bowel Preparation Reduces Incisional Surgical Site Infection and Anastomotic Leak Rates After Elective Colorectal Resection: An Analysis of Colectomy-Targeted ACS NSQIP. *Ann Surg.* 2015;262(2):331-7.
94. Kiran RP, Murray AC, Chiuzan C, Estrada D, Forde K. Combined pre-operative mechanical bowel preparation with oral antibiotics significantly reduces surgical site infection, anastomotic leak, and ileus after colorectal surgery. *Ann Surg.* 2015;262(3):416-25; discussion 23-5.

95. Moghadamyeghaneh Z, Hanna MH, Carmichael JC, Mills SD, Pigazzi A, Nguyen NT, et al. Nationwide Analysis of Outcomes of Bowel Preparation in Colon Surgery. *J Am Coll Surg*. 2015;220(5):912-20.
96. ClinicalTrials.gov. Does Administration of Antibiotics in Patients Undergoing Surgery for Colorectal Cancer Result in Less Complications and Better Prognosis? (SELECT) 2013 [05/02/2016]. Available from: <https://clinicaltrials.gov/ct2/show/NCT01740947>.
97. den Dulk M NS, Hendriks ER, et al. Improved diagnosis and treatment of anastomotic leakage after colorectal surgery. *Eur J Surg Oncol*. 2009;35:7.
98. Essani RB BR. Anastomotic leak in colorectal surgery: a review. *Gastroenterol Polska*. 2009;16:5.
99. Shorthouse AJ BC, Evers AA, Thompson JPS. The water soluble contrast enema after rectal anastomosis. *Br j Surg*. 1982;69:4.
100. Akyol AM MJ, Galloway DJ, George WD. Early postoperative contrast radiology in the assessment of colorectal anastomotic integrity. *Int J Colorectal Dis*. 1992;7:3.
101. Nesbakken A NK, Lunde OC, Blucher J, Gjertsen O, Dullerud R. Anastomotic leak following mesorectal excision for rectal cancer: true incidence and diagnostic challenges. *Colorect Dis*. 2005;7:5.
102. Bertoni CB MM, Fleury AR, VanderMeer TJ, Skeist BP, Cagir B. Utility of pelvic CT with rectal contrast to identify pelvic abscess and anastomotic leaks. *Gastroenterol*. 2009;136.
103. Millan M, Garcia-Granero E, Flor B, Garcia-Botello S, Lledo S. Early prediction of anastomotic leak in colorectal cancer surgery by intramucosal pH. *Dis Colon Rectum*. 2006;49(5):595-601.
104. Ellebaek Pedersen M, Qvist N, Bisgaard C, Kelly U, Bernhard A, Moller Pedersen S. Peritoneal microdialysis. Early diagnosis of anastomotic leakage after low anterior resection for rectosigmoid cancer. *Scand J Surg*. 2009;98(3):148-54.
105. Miller K, Arrer E, Leitner C. Early detection of anastomotic leaks after low anterior resection of the rectum. *Dis Colon Rectum*. 1996;39(10):1081-5.
106. Junger W, Junger WG, Miller K, Bahrami S, Redl H, Schlag G, et al. Early detection of anastomotic leaks after colorectal surgery by measuring endotoxin in the drainage fluid. *Hepatogastroenterology*. 1996;43(12):1523-9.
107. Fouda E, El Nakeeb A, Magdy A, Hammad EA, Othman G, Farid M. Early detection of anastomotic leakage after elective low anterior resection. *J Gastrointest Surg*. 2011;15(1):137-44.
108. DeArmond DT CA, Johnson SB. Anastomotic leak detection by electrolyte electrical resistance. *J Invest Surg*. 2010;23:12.
109. Daams F, Wu Z, Lahaye MJ, Jeekel J, Lange JF. Prediction and diagnosis of colorectal anastomotic leakage: A systematic review of literature. *World Journal of Gastrointestinal Surgery*. 2014;6(2):14-26.
110. Daams F, Sliker JC, Tedja A, Karsten TM, Lange JF. Treatment of colorectal anastomotic leakage: results of a questionnaire amongst members of the Dutch Society of Gastrointestinal Surgery. *Dig Surg*. 2012;29(6):516-21.
111. Moghadamyeghaneh Z, Hanna MH, Alizadeh RF, Carmichael JC, Mills S, Pigazzi A, et al. Contemporary management of anastomotic leak after colon surgery: assessing the need for reoperation. *The American Journal of Surgery*. 2015.
112. Rickert A, Willeke F, Kienle P, Post S. Management and outcome of anastomotic leakage after colonic surgery. *Colorectal Dis*. 2010;12(10):e216-23.

113. Weidenhagen R, Gruetzner KU, Wiecken T, Spelsberg F, Jauch K-W. Endoscopic vacuum-assisted closure of anastomotic leakage following anterior resection of the rectum: a new method. *Surgical Endoscopy*. 2008;22(8):1818-25.
114. van Koperen PJ, van Berge Henegouwen MI, Rosman C, Bakker CM, Heres P, Slors JF, et al. The Dutch multicenter experience of the endo-sponge treatment for anastomotic leakage after colorectal surgery. *Surg Endosc*. 2009;23(6):1379-83.
115. Weiland T, Fehlker M, Gottwald T, Schurr MO. Performance of the OTSC System in the endoscopic closure of iatrogenic gastrointestinal perforations: a systematic review. *Surg Endosc*. 2013;27(7):2258-74.